

DOLFIN



Supplement Dosing Charts

TABLE 1: GIVING THE SUPPLEMENT AS A SHOT: BEFORE 1 FEED PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you give the supplement before 1 feed per day

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	If given in 1 feed per day, number of scoops of supplement per feed	Minimum volume of milk for mixing supplement (ml)
	Weight between ... and ... kg			
1	1.0-1.99	1	1	3
2	2.0-2.99	2	2	6
3	3.0-3.99	3	3	9
4	4.0-4.99	4	4	12
5	5.0-5.99	5	5	15
6	6.0-6.99	6	6	18
7	7.0-7.99	7	7	21
8	8.0-8.99	8	8	24
9	9.0-9.99	9	9	27
10	10.0-10.99	10	10	30
11	11.0-11.99	11	11	33
12+	12 kg or more	12	12	36

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 2: GIVING THE SUPPLEMENT AS A SHOT: BEFORE 2 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you give the supplement before 2 feeds per day

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	Number of scoops with supplement per feed (if spread across 2 feeds per day)		Minimum volume of milk for mixing supplement (ml)	
			Feed 1	Feed 2	Feed 1	Feed 2
	Weight between ... and ... kg					
2	2.0-2.99	2	1	1	3	3
3	3.0-3.99	3	2	1	6	3
4	4.0-4.99	4	2	2	6	6
5	5.0-5.99	5	3	2	9	6
6	6.0-6.99	6	3	3	9	9
7	7.0-7.99	7	4	3	12	9
8	8.0-8.99	8	4	4	12	12
9	9.0-9.99	9	5	4	15	12
10	10.0-10.99	10	5	5	15	15
11	11.0-11.99	11	6	5	18	15
12+	12 kg or more	12	6	6	18	18

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 3: GIVING THE SUPPLEMENT AS A SHOT: BEFORE 3 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you give the supplement before 3 feeds per day

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	Number of scoops with supplement per feed (if spread across 3 feeds per day)			Minimum volume of milk for mixing supplement (ml)		
			Feed 1	Feed 2	Feed 3	Feed 1	Feed 2	Feed 3
3	3.0-3.99	3	1	1	1	3	3	3
4	4.0-4.99	4	2	1	1	6	3	3
5	5.0-5.99	5	2	2	1	6	6	3
6	6.0-6.99	6	3	2	1	9	6	3
7	7.0-7.99	7	3	2	2	9	6	6
8	8.0-8.99	8	3	3	2	9	9	6
9	9.0-9.99	9	3	3	3	9	9	9
10	10.0-10.99	10	4	4	2	12	12	6
11	11.0-11.99	11	4	4	3	12	12	9
12+	12 kg or more	12	4	4	4	12	12	12

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 4: GIVING THE SUPPLEMENT AS A SHOT: BEFORE 4 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you give the supplement before **4 feeds per day**

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	Number of scoops with supplement per feed (if spread across 4 feeds per day)				Minimum volume of milk for mixing supplement (ml)			
			Feed 1	Feed 2	Feed 3	Feed 4	Feed 1	Feed 2	Feed 3	Feed 4
	Weight between ... and ... kg									
4	4.0-4.99	4	1	1	1	1	3	3	3	3
5	5.0-5.99	5	2	1	1	1	6	3	3	3
6	6.0-6.99	6	2	2	1	1	6	6	3	3
7	7.0-7.99	7	3	2	1	1	9	6	3	3
8	8.0-8.99	8	3	2	2	1	9	6	6	3
9	9.0-9.99	9	3	2	2	2	9	6	6	6
10	10.0-10.99	10	3	3	2	2	9	9	6	6
11	11.0-11.99	11	3	3	3	2	9	9	9	6
12+	12 kg or more	12	3	3	3	3	9	9	9	9

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

Giving the supplement by bottle or tube feeding

The following tables show the amount of supplement needed per day and per feed based on the weight of your child when given by bottle or tube feeding with 1, 2, 3 or 4 feeds.

TABLE 5: GIVING THE SUPPLEMENT BY BOTTLE OR BY TUBE FEEDING: IN 1 FEED PER DAY

*Amount of supplement per day, based on the weight of your child, if you give the supplement in **1 feed per day***

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	If given in 1 feed per day, number of scoops of supplement per feed	Minimum volume of milk for mixing supplement (ml)
	Weight between ... and ... kg			
1	1.0-1.99	1	1	15
2	2.0-2.99	2	2	30
3	3.0-3.99	3	3	45
4	4.0-4.99	4	4	60
5	5.0-5.99	5	5	75
6	6.0-6.99	6	6	90
7	7.0-7.99	7	7	105
8	8.0-8.99	8	8	120
9	9.0-9.99	9	9	135
10	10.0-10.99	10	10	150
11	11.0-11.99	11	11	165
12+	12 kg or more	12	12	180

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 6: GIVING THE SUPPLEMENT BY BOTTLE OR BY TUBE FEEDING: IN 2 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you spread the supplement across 2 feeds per day

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	Number of scoops with supplement per feed (if spread across 2 feeds per day)		Minimum volume of milk for mixing supplement (ml)	
			Feed 1	Feed 2	Feed 1	Feed 2
	Weight between ... and ... kg					
2	2.0-2.99	2	1	1	15	15
3	3.0-3.99	3	2	1	30	15
4	4.0-4.99	4	2	2	30	30
5	5.0-5.99	5	3	2	45	30
6	6.0-6.99	6	3	3	45	45
7	7.0-7.99	7	4	3	60	45
8	8.0-8.99	8	4	4	60	60
9	9.0-9.99	9	5	4	75	60
10	10.0-10.99	10	5	5	75	75
11	11.0-11.99	11	6	5	90	75
12+	12 kg or more	12	6	6	90	90

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 7: GIVING THE SUPPLEMENT BY BOTTLE OR BY TUBE FEEDING: IN 3 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you give the supplement across 3 feeds per day

Weight category	Child's weight (in kg)	Number of scoops of supplement per day	Number of scoops with supplement per feed (if spread across 3 feeds per day)			Minimum volume of milk for mixing supplement (ml)		
			Feed 1	Feed 2	Feed 3	Feed 1	Feed 2	Feed 3
	Weight between ... and ... kg							
2	2.0-2.99	2	1	1	None	15	15	None
3	3.0-3.99	3	1	1	1	15	15	15
4	4.0-4.99	4	2	1	1	30	15	15
5	5.0-5.99	5	2	2	1	30	30	15
6	6.0-6.99	6	3	2	1	45	30	15
7	7.0-7.99	7	3	2	2	45	30	30
8	8.0-8.99	8	3	3	2	45	45	30
9	9.0-9.99	9	3	3	3	45	45	45
10	10.0-10.99	10	4	4	2	60	60	30
11	11.0-11.99	11	4	4	3	60	60	45
12+	12 kg or more	12	4	4	4	60	60	60

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

TABLE 8: GIVING THE SUPPLEMENT BY BOTTLE OR BY TUBE FEEDING: IN 4 FEEDS PER DAY

Amount of supplement needed per day and per feed based on the weight of your child, if you spread the supplement across 4 feeds per day

Weight category	Child's weight (in kg)	Number of scoops with supplement per day	Number of scoops with supplement per feed (if spread across 4 feeds per day)				Minimum volume of milk for mixing supplement (ml)			
			Feed 1	Feed 2	Feed 3	Feed 4	Feed 1	Feed 2	Feed 3	Feed 4
2	2.0-2.99	2	1	1	None	None	15	15	None	None
3	3.0-3.99	3	1	1	1	None	15	15	15	None
4	4.0-4.99	4	1	1	1	1	15	15	15	15
5	5.0-5.99	5	2	1	1	1	30	15	15	15
6	6.0-6.99	6	2	2	1	1	30	30	15	15
7	7.0-7.99	7	3	2	1	1	45	30	15	15
8	8.0-8.99	8	3	2	2	1	45	30	30	15
9	9.0-9.99	9	3	2	2	2	45	30	30	30
10	10.0-10.99	10	3	3	2	2	45	45	30	30
11	11.0-11.99	11	3	3	3	2	45	45	45	30
12+	12 kg or more	12	3	3	3	3	45	45	45	45

General rule: 1 scoop of supplement needs to be dissolved in at least 15 ml expressed breast milk or infant formula.

Contact address:

NPEU Clinical Trials Unit
National Perinatal Epidemiology Unit (NPEU)
Nuffield Department of Population Health
University of Oxford, Old Road Campus, Oxford OX3 7LF

T: 01865 617919

E: dolphin@npeu.ox.ac.uk

W: www.npeu.ox.ac.uk/dolphin

This study is being organised by the National Perinatal Epidemiology Unit (NPEU) CTU at the University of Oxford. The Unit is dedicated to improving the care provided to women and their families during pregnancy, childbirth and the period after birth, as well as the care provided to the newborn. www.npeu.ox.ac.uk

REC Reference: 22/SW/0009 ISRCTN: 62323236

This project was funded by the National Institute for Health and Care Research Health Technology Assessment (HTA) Programme. (Project number NIHR 130925).